



King's Research Portal

DOI:

[10.1073/pnas.1502902112](https://doi.org/10.1073/pnas.1502902112)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Wefelmeyer, W., Cattaert, D., & Burrone, J. (2015). Activity-dependent mismatch between axo-axonic synapses and the axon initial segment controls neuronal output. *Proceedings of the National Academy of Sciences of the United States of America*. <https://doi.org/10.1073/pnas.1502902112>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Activity-dependent mismatch between axo-axonic synapses and the axon initial segment controls neuronal output

Winnie Wefelmeyer^a, Daniel Cattaert^b, and Juan Burrone^{a,1}

^aMRC Centre for Developmental Neurobiology, King's College London, London SE1 1UL, United Kingdom; and ^bInstitut de Neurosciences Cognitives et Intégratives d'Aquitaine, Université de Bordeaux, CNRS, 33076 Bordeaux, France

Edited by Gina G. Turrigiano, Brandeis University, Waltham, MA, and approved June 26, 2015 (received for review February 11, 2015)

The axon initial segment (AIS) is a structure at the start of the axon with a high density of sodium and potassium channels that defines the site of action potential generation. It has recently been shown that this structure is plastic and can change its position along the axon, as well as its length, in a homeostatic manner. Chronic activity-deprivation paradigms in a chick auditory nucleus lead to a lengthening of the AIS and an increase in neuronal excitability. On the other hand, a long-term increase in activity in dissociated rat hippocampal neurons results in an outward movement of the AIS and a decrease in the cell's excitability. Here, we investigated whether the AIS is capable of undergoing structural plasticity in rat hippocampal organotypic slices, which retain the diversity of neuronal cell types present at postnatal ages, including chandelier cells. These interneurons exclusively target the AIS of pyramidal neurons and form rows of presynaptic boutons along them. Stimulating individual CA1 pyramidal neurons that express channelrhodopsin-2 for 48 h leads to an outward shift of the AIS. Intriguingly, both the pre- and postsynaptic components of the axo-axonic synapses did not change position after AIS relocation. We used computational modeling to explore the functional consequences of this partial mismatch and found that it allows the GABAergic synapses to strongly oppose action potential generation, and thus downregulate pyramidal cell excitability. We propose that this spatial arrangement is the optimal configuration for a homeostatic response to long-term stimulation.

axon initial segment | chandelier cells | optogenetics | intrinsic plasticity | homeostatic plasticity

Neurons receive a large number of synaptic inputs along the somato-dendritic compartment that integrate at the axon initial segment (AIS) to fire an action potential (AP) (1–3). As an important site for transforming graded synaptic inputs into all-or-none APs, it is also a potentially sensitive target for the modulation of neuronal excitability (4, 5). In fact, one interesting aspect of principal neurons in the hippocampus and cortex is the presence of a unique type of GABAergic axo-axonic synapse that forms onto the AIS and controls neuronal output (6, 7). These synapses are formed by a specific group of fast-spiking interneurons, the chandelier cells, that are generally found sparsely distributed in the brain and have therefore been difficult to study in the past (6, 8). However, the recently developed transgenic mouse lines that can label chandelier neurons more selectively have begun to shine light on their form and function (9). In the cortex, the axonal “cartridges” of synaptic boutons that form onto the AIS are generally found on the more distal AIS domain, where the AP is thought to initiate (9). Although there is little information on the role of these interneurons in network function, they have been implicated in a number of events, including driving negative feedback in the dentate gyrus (10), modulating the emergence of sharp waves in the hippocampus (11), providing complex inhibitory-excitatory feedforward loops in the cortex (12), as well as shunting and filtering out APs back-propagating from ectopic initiation sites in the hippocampus (13). In addition, there is an ongoing debate as to whether axo-

axonic synapses depolarize or hyperpolarize postsynaptic pyramidal neurons (14), although in conditions mimicking *in vivo*-like oscillations, they act mainly as inhibitors of excitability (15). The emerging picture is one where chandelier cells play an important role in modulating neuronal output at the AIS, but the precise way in which this happens remains to be properly established.

More recently, longer-term forms of modulation have been described at the AIS in response to chronic alterations in neuronal activity (16–20). For example, sensory deprivation of chick brainstem auditory neurons caused an increase in the length of the AIS, which was paralleled by an increase in neuronal excitability (18). Conversely, in dissociated hippocampal neurons, increases in neuronal activity through either application of hyperkalemic solution or by optogenetic means resulted in the distal relocation of the AIS, which correlated with a decrease in excitability (16, 17). These exciting forms of structural and functional plasticity have raised many questions that remain unanswered. For example, what happens to axo-axonic synapses when the AIS relocates? Do they also relocate to remain in register with the AIS or do they remain static, causing a mismatch between the two? Implicit in this question is the idea that different spatial arrangements between the AIS and axo-axonic synapses (eg: “in register” versus “mismatched”) could have very different consequences on neuronal output.

In this study we performed optogenetic activation of individual CA1 pyramidal neurons and show that activity-dependent relocation of the AIS occurs in hippocampal slices. However, the axo-axonic synapses that form on the AIS do not change their position, resulting in a mismatch between the two compartments. By using computational modeling of a typical CA1 neuron, we find that the distal relocation of the AIS, together with the activation of axo-axonic synapses in the proximal axonal domain, is

Significance

Neurons integrate synaptic inputs at the axon initial segment (AIS), where the action potential is initiated. Recent findings have shown this structure to be highly plastic. Here, we focus on the axo-axonic synapses formed onto the AIS and show that although chronic increases in neuronal activity result in a distal relocation of the AIS, the synapses do not change position. Computational modeling suggests this spatial mismatch has critical functional consequences on neuronal output, allowing the cell to downregulate its own excitability and thus respond homeostatically to chronic stimulation.

Author contributions: W.W. and J.B. designed research; W.W. performed research; W.W. analyzed data; W.W. and J.B. wrote the paper; and D.C. provided the mathematical model.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

¹To whom correspondence should be addressed. Email: juan.burrone@kcl.ac.uk.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1502902112/-DCSupplemental.

the perfect combination for decreasing neuronal excitability. Our findings suggest that the interplay between the position of the AIS and its synapses is another important player in the set of homeostatic responses used by neurons to adapt to chronic changes in activity.

Results

In principal neurons of the hippocampus and cortex, the AIS receives GABAergic synaptic modulation through axo-axonic synapses formed by chandelier interneurons (6, 21). We set out to label the AIS and its corresponding axo-axonic synapses in CA1 and CA3 pyramidal neurons from organotypic slices of the hippocampus. Staining for ankyrin-G (AnkG) resulted in a strong label of the AIS (Fig. 1). Axo-axonic synapses were colabeled with antibodies against either gephyrin, the postsynaptic structural marker for GABAergic synapses, or against the $\alpha 2$ subunit of GABA_A receptors (GABA_A $\alpha 2$), which is preferentially targeted to

synapses along the AIS (22). Labeling of presynaptic GABAergic boutons by staining for the vesicular GABA transporter, VGAT, resulted in a strong colocalization with both postsynaptic markers. Close examination of the images showed that although AIS length was comparable between CA1 and CA3 neurons, the density and overall number of axo-axonic synapses formed onto the AIS of CA3 neurons was larger than that of CA1 cells (Fig. 1*B* and *C*). The spatial distribution of synapses along the AIS was similar in both neurons and, interestingly, many synapses were found beyond the AIS (19, 23). This finding was further confirmed by imaging chandelier cell axons forming axo-axonic synapses along and beyond the AIS of CA1 neurons (Fig. 1*F*). One striking observation that was apparent in the confocal images of the thick CA3 axons was the heterogeneous distribution of the AnkG stain, showing gaps or holes along the axon. More importantly, we found that these AnkG-empty domains were often filled (or partially filled) with staining for postsynaptic markers of axo-axonic synapses (Fig. 1*D* and *E*), as previously described (24). Although we did not see a similar staining pattern in the much thinner CA1 neurons, previous work using superresolution microscopy did show a separation of AIS and synaptic compartments, suggesting that this is a general feature of pyramidal neurons in the hippocampus and neocortex (24). These findings suggest the existence of two separate domains within the AIS, governed by different scaffolding proteins that form nonoverlapping macromolecular complexes and may therefore be controlled independently of each other. Because the AIS has been shown to undergo an activity-dependent relocation of its AnkG-based domain in dissociated hippocampal neurons (16, 17, 19), we sought to establish whether axo-axonic synapses also relocated, to remain in register with the AIS.

Single-Cell Optogenetic Stimulation Induces AIS Relocation in Organotypic Hippocampal Slices

We performed chronic stimulation of individual CA1 pyramidal neurons expressing channelrhodopsin-2 (ChR2), using a previously described optogenetic approach (17). As controls, we used EGFP-expressing neurons from sister slices, grown in the same incubator, exposed to the same amount of blue LED light. After fixing and staining for AnkG, the spatial distribution of the AIS was measured in 3D (Figs. 1*A* and 2*A* and *B*). One immediate observation from these images was that CA1 neurons could be classified into two broad morphological groups: those with an AIS that rises from the soma and those that branch off from a basal dendrite (25). Regardless of the morphology of the neuron, stimulated neurons showed a change in the position of the AIS such that the start, maximum, and end position of the AnkG stain was significantly shifted along the axon by about 12 μ m. This finding was also true when measurements were taken from the axon origin, be it a basal dendrite or the soma. The length of the AIS did not change. Previous findings (17) have linked changes in AIS structure to changes in neuronal excitability. Indeed, whole-cell patch-clamp recordings revealed a significantly higher current threshold and a dramatic rightward shift of input-output curves in stimulated neurons (Fig. S1), indicating a decrease in neuronal excitability. In parallel to this, the input resistance (R_{in}) decreased significantly, which could not be explained by changes in hyperpolarization-activated currents (I_h) or barium-sensitive potassium conductances, but is caused, in large part, by a change in tonic GABA_AR conductance (Fig. S2). Throughout our experiments we have used gabazine (SR95531; 10 μ M) to block GABA_A receptors, which only has a limited effect on tonic GABA_A conductances (26–28). Blocking both synaptic and tonic GABA_AR conductances with Picrotoxin (100 μ M) lead to a significant reduction in threshold current and a large leftward shift of the input-output curve of ChR2-stimulated neurons (Fig. S3). This finding suggests that a tonic GABA_AR conductance is regulated in an activity-dependent





Fig. 1. Axo-axonic synapses in organotypic hippocampal slices. (A) A ChR2-transfected CA3 pyramidal neuron, stained with AnkG (magnification). Start, maximum, and end position of the AIS was determined using the fluorescent intensity of the AnkG label along the axonal process. (Scale bar, Left: 20 μ m.) (B) VGAT and AnkG staining in the CA3 and CA1 regions of the hippocampus. (C) AIS length and number of VGAT puncta on AISs in the CA3 and CA1 regions (means \pm SEM; * P < 0.05; n = 16 CA1; n = 17 for CA3 AIS length; n = 12 for CA3 puncta). (D) GABA $\alpha 2$ and AnkG staining in the CA3 region. Arrows indicate apparent holes in the AnkG staining, filled with GABA $\alpha 2$ puncta. (E) Magnification of boxed areas as shown in *D*. Fluorescent intensity traces were taken through AIS and puncta as indicated by the white lines. (Scale bars, 20 μ m.) (F) CA3 pyramidal neuron expressing ChR2-YFP in Nkx2.1^{CreER::Ai9} mouse showing chandelier cell boutons along its axon (arrows). AIS position is marked with arrowheads (Right). Dotted vertical line indicates image concatenation.



manner and can account for much of the drop in neuronal output.

Axo-Axonic Synapses Do Not Change Position After AIS Plasticity. Do axo-axonic synapses that form onto the AIS of CA1 pyramidal neurons also relocate after chronic stimulation? Stimulated neurons were stained for AnkG and the synaptic markers VGAT and GABA_Aα2. We found that although the AIS did relocate, the GABAergic synapses did not change position (Fig. 2 *E–H*) (19). The distribution of VGAT and GABA_Aα2 terminals along the axon was identical to control neurons, suggesting that axo-axonic synapses are controlled independently of the AIS. A distribution of synapse position relative to the start of the AIS shows a shift in stimulated neurons compared with controls, with synapses being left behind in the space between the start of the

AIS and the soma. Importantly, the distal relocation of the AIS did not lead to a major change in the number of synapses that form along it, because GABAergic synapses are also found beyond the AIS in unstimulated neurons (Fig. 1*F* and Fig. S4) (19, 23). We also saw no difference in the colocalization of pre- and postsynaptic markers, precluding the possibility that postsynaptic compartments moved with the AIS leaving orphan presynaptic boutons behind (or vice versa).

An Optimal Spatial Arrangement of the AIS and Its Synapses for Downregulating Neuronal Activity. To establish whether this mismatch between axo-axonic synapses and the AIS has a functional role, we used a simple model of a CA1 neuron that contained a well-defined AIS and axo-axonic synapses at the same density to that observed in our experiments (Fig. 34) (29). Our model




Fig. 3. Computational model examining the effect of axo-axonic synapse position on APs. (A) GABAergic synapses were evenly spaced along the axon starting at 2 μm from the soma in the control and “AIS distal” scenario. In the third scenario the synapses were displaced distally together with the AIS. (B) APs elicited by a current pulse in the three scenarios. (C) Peak AP voltage (Left) and difference in peak voltage compared with control (Right), for varying GABA_AR conductance. (D) Time to AP peak (Left) and difference in time to peak compared with control (Right). (E) Current threshold in response to a short, 3-ms current injection (Left) and difference in current threshold compared with control (Right).

compared a control neuron with the two conceivable arrangements after AIS relocation: (*i*) a neuron with a distal AIS and proximal synapses, resulting in a mismatch as observed experimentally; and (*ii*) a neuron with a distal AIS and distal synapses, so that both remain in register with each other. Activating GABA_ARs in these three conditions revealed important functional consequences on neuronal output and highlights the critical importance of synapse location relative to the AIS. Neurons with a distal AIS and proximal synapses (mismatched) showed delayed APs that were smaller in amplitude compared with controls. On the other hand, neurons with a distal AIS and distal synapses (in register) showed a shorter lag to the first AP and a change in amplitude that was highly dependent on the levels of GABA_AR activation (Fig. 3 *C* and *D*). Importantly, the delays in AP firing were within the time-domain (tens of milliseconds) important for spike-timing-dependent plasticity rules that, when combined with changes in AP amplitude, will have

important consequences for long-term forms of synaptic plasticity. Finally, we also found that neurons with a distal AIS, but proximal synapses had a higher current threshold, whereas those with distal synapses actually showed a lower current threshold compared with controls (Fig. 3*E*).

Discussion

The findings presented here show that optogenetic activation of CA1 pyramidal neurons in the hippocampus resulted in a distal relocation of the AIS, but no change in the position of axo-axonic synapses, creating a mismatch between the AIS and its GABAergic inputs. More importantly, our model predicts that this configuration has multiple and far-reaching effects on AP initiation and properties, including modulation of the timing and amplitude of back-propagating APs, both of which are likely to affect long-term plasticity of synaptic inputs. The current threshold to short stimuli is also increased, making it more efficient at filtering out depolarizing synaptic inputs, thus resulting in a homeostatic decrease in neuronal excitability. Interestingly, our model predicted that a relocation of the synapses together with the AIS would have had the opposite effect, an increase in neuronal excitability compared with controls. It appears that the dissociation between the AIS and its axo-axonic synapses leads to an optimal configuration for a homeostatic response to long-term stimulation. Our findings reveal a novel form of plasticity where the position of synapses and their target can be modified with exquisite precision to fine-tune neuronal output.

Distribution of Axo-Axonic GABAergic Synapses in the Hippocampus.

An activity-dependent change in AIS position has previously been reported in dissociated hippocampal neurons (17). Here we show, to our knowledge for the first time, that a similar form of plasticity also occurs in CA1 neurons in organotypic slices, a preparation where the architecture of the network and its connections is preserved. In addition, slices can be taken from post-natal tissue, assuring that the late-born chandelier cells have migrated to the hippocampus before slice preparation (9). As a result, we were able to visualize the GABAergic axo-axonic synapses made by chandelier neurons onto the AIS of pyramidal cells and follow their behavior during AIS plasticity. Axo-axonic synapses formed all along the AIS and were not biased toward the distal end, as observed in cortical neurons. Strikingly, we also observed that synapses were not spatially constrained to the AIS but extended beyond it (19, 23). Although it is unclear what role these synapses play, it is likely that they modulate AP propagation down the axon. In the context of AIS plasticity, these synapses ensure that the AIS is still innervated even after it relocates distally, so that axo-axonic synapses can continue to modulate neuronal output.

Along the axon of CA3 neurons, axo-axonic synapses often fitted in the gaps or holes in the AnkG stain of the AIS. This punctured distribution of AnkG has been previously described in a number of different neurons, including in CA1 and CA3 cells, using superresolution structured illumination microscopy (24). The gaps in AnkG were filled mainly by clusters of the voltage-gated potassium channel Kv2.1, the cisternal organelle marker synaptopodin, and GABAergic synapses. Further high-resolution imaging will need to be done following activity-dependent AIS relocation, especially on the thin axons of CA1 pyramidal cells, to establish what happens to the ultrastructure of the AIS, its synapses, and other proteins associated with axo-axonic synapses. The Kv2.1 channel is particularly interesting because it does not have an AnkG binding domain (unlike other ion channels at the AIS) and undergoes an activity-dependent dispersion (loss of clustering) and change in gating properties thought to be homeostatic in nature (30, 31). These events are calcineurin-dependent (30), much like the AIS relocation in dissociated neurons (16), implying that similar signaling cascades may be acting on both

